
Influenza and Related Disease Updates for Los Angeles County

Updated: 22NOV2019 MMWR week: 46 Ending on: 11/16/2019 Season: 2019-2020

This week at a glance

- Indicators of influenza activity in Los Angeles county have increased over the past two weeks.
- More influenza B has been detected than influenza A in Los Angeles County over the past two weeks. It's still too early to know which strains will predominate this season.
- It's not too late to get vaccinated! Vaccination is the best way to prevent flu and its potentially serious complications.



_				
ED-IL	This week	32	Rate per 1,000 visits to a Los Angeles County* emergency department were for influenza-like illness	♠
	Last week	28	(ED-ILI).	

P&I	This week	7.8%	Number of <i>deaths</i> recorded in Los Angeles County*	
	Last week	7.1%	were caused by <i>pneumonia or influenza</i> (P&I).	

ths			
ភ	Season to date**	2	Number of <i>influenza-associated deaths</i> have been reported to Los Angeles County DPH.

*Los Angeles County DPH surveillance data excludes the cities of Long Beach and Pasadena.

**The influenza-surveillance period starts with MMWR week 40 and runs through week 39 of the following year. The 2019-20 season started on September 29, 2019.


Los Angeles County Department of Public Health (DPH) prepares this newsletter to summarize current influenza surveillance data in Los Angeles County. *Weekly surveillance data are preliminary and subject to change.* More information regarding methods can be found on the surveillance system specific pages of this report.

Influenza and Related Disease Updates for Los Angeles County

Updated: 22NOV2019 MMWR week: 46 Ending on: 11/16/2019 Season: 2019-2020


Sentinel Laboratory Surveillance


Viral surveillance data is provided by eight clinical laboratories serving hospitals and healthcare networks across Los Angeles County. Participating laboratories provide the number of positive tests and total number of specimens tested for influenza and respiratory syncytial virus. Data is reported on a weekly basis; weeks start on Sunday and end the following Saturday. Many participating laboratories also report data on other respiratory viruses.




Respiratory specimens testing positive for Influenza at LAC Sentinel

100 5% 0 0% 40 42 44 46 48 50 52 1 3 5 7 9 11 13 15 17 19 21 23 25 27 29 31 33 35 37 39 MMWR Week








Influenza and Related Disease Updates for Los Angeles County


Influenza-like illness

ACDC's Syndromic Surveillance Project monitors initial self-reported symptoms from patients presenting to participating emergency departments throughout Los Angeles County. These symptoms are categorized into different clinical syndromes according to specific code words. The syndrome of ILI includes symptoms such as: fever, congestion, sneezing, sore throat, runny nose, and cough. The proportion of ILI emergency department (ED) visits for all ages and by age group is analyzed weekly and is conducted year-round.

Emergency department visits for influenza-like illness, per 1,000, Los Angeles County 2015-16 through 2019-20 influenza seasons



Emergency department visits for influenza-like illness by age groups, Los Angeles County, 2019-20 influenza season




Influenza and Related Disease Updates for Los Angeles County

Updated: 22NOV2019 MMWR week: 46 Ending on: 11/16/2019 Season: 2019-2020

Pneumonia and Influenza

Each week, the Office of Health Assessment and Epidemiology at LACDPH reports the total number of death certificates received and the number of those for which pneumonia or influenza (P&I) was listed as the underlying or contributing cause of death by age group. The percentage of all deaths due to P & I is calculated and compared to previous season's data.



Vital statistic data :*The reporting period is a 7-day week ending on Saturday. The number of deaths is the number of certificates received by the Office of Vital Records during the reporting period regardless of date of death. The count includes all certificates of deaths (excludes fetal deaths) occurring in the County of Los Angeles* regardless of the residence of the deceased. As a result, many deaths included in this report are from out of county.*Deaths that occurred in Long Beach and Pasadena are excluded. When confirmed, any influenza associated deaths are included in the death count above.

Pneumonia and Influenza (P&I) Listed As a Cause of Death, Number and Percent of All Deaths Los Angeles County, 2019-20

Influenza and Related Disease Updates for Los Angeles County

 Updated:
 22NOV2019

 MMWR week:
 46

 Ending on:
 11/16/2019

 Season:
 2019-2020

Deaths

Deaths associated with influenza infection are reportable in all ages in Los Angeles county. A death is confirmed as being influenza-associated when there is a laboratory confirmed influenza infection, the cause of death is clinically compatible with influenza or influenza complications, and there was no return to baseline health between infection and death. Clinically compatible complications can include pneumonia and cardiovascular problems like heart attacks. Influenza can also exacerbate long-term medical conditions (such as COPD, heart failure, or diabetes) which can lead to death.

	20	19-20	20:	18-19	20	17-18	20	16-17	20	15-16	20:	14-15
Confirmed deaths		2	1	L 25		285		86		77		55
Mean age (STD)	52.5	(13.4)	67.0	(18.9)	75.4	(16.3)	77.0	(17.7)	61.3	(23.3)	70.2	(26.6)
Range (Low-High)			1	- 104	9	- 105	4	- 102	1	- 103	1	- 101
Median age (Q1, Q3)			68	(56 ,81)	79	(66 ,87)	82.5	(66 ,89)	61	(47 ,80)	81	(55 ,90)
	n	%	n	%	n	%	n	%	n	%	n	%
Age												
0-5 years	0		2	(1.6)	0		1	(1.2)	2	(2.6)	1	(1.8)
6 - 17 years	0		0		2	(.7)	0		1	(1.3)	2	(3.6)
18-64 years	2	(100.0)	46	(36.8)	62	(21.8)	20	(23.3)	39	(50.6)	14	(25.5)
65+ years	0		77	(61.6)	221	(77.5)	65	(75.6)	35	(45.5)	38	(69.1)

The number of influenza-associated deaths reported to LAC DPH does not represent the true mortality associated with influenza in Los Angeles County. Public health authorities recognize that current surveillance methods substantially undercount influenza-associated deaths. Most people who get influenza do not seek care. Most people who seek care are not tested for influenza. Severe complications of influenza may occur after the virus is no longer detectable in the body. Testing practices may vary across seasons. For these reasons, between season comparison of reported influenza-associated deaths may not be reliable.

Outbreaks

Outbreaks of any disease are reportable in Los Angeles County. Respiratory outbreak definitions vary by setting; in general, the occurrence of a cluster of influenza-like illness (Fever>100 F with cough and/or sore throat) is cause for investigation. Suspect outbreaks are currently under investigation.

		Season	to date Previous seasons					
		Confirmed Suspect		2018-19 2017-18		2016-17 2015-16		2014-15
Health Facilities	Influenza	0	0	18	70	29	11	24
	Other	0	0	0	0	0	0	0
Facilities	Unknown	0	0	4	12	2	5	7
	Influenza	0	0	7	37	6	10	18
Community	Other	0	0	0	0	0	0	0
	Unknown	0	1	19	35	27	20	27
Total		0	1	48	154	64	46	76

Los Angeles Couty Department of Public Health Acute Communicable Disease Control (213) 240-7941 www.publichealth.lacounty.gov/acd