
Congenital Syphilis




Congenital Syphilis California Incidence Rates, 1940–2017






Congenital Syphilis, California versus United States Incidence Rates, 1963–2017






Note: The Modified Kaufman Criteria were used through 1989. The CDC Case
Definition (MMWR 1989; 48: 828) was used effective January 1, 1990.
California data prior to 1985 include all cases of congenital syphilis, regardless of age.

Congenital Syphilis Incidence Rates by County, California, 2017




Congenital Syphilis Number of Cases by County, California, 2017





Ranking of County Congenital Syphilis Rates California, 2017

(with 95% Confidence Intervals*)






Note: Rates are per 100,000 live births.

Source: California Department of Public Health, STD Control Branch

^{*} Confidence intervals were calculated using Poisson exact method; excludes counties with no cases.


Congenital Syphilis Cases by Vital Status and Presence of Signs/Symptoms California, 2008–2017






Note: Alive, no signs/symptoms includes alive with missing documentation on signs/symptoms. Of the 283 total cases, 233 alive with no signs, 14 alive with signs of CS, 36 stillbirth or neonatal death

Congenital Syphilis Stillbirths California, 2011–2017






Congenital Syphilis California, 1990–2017





Congenital Syphilis versus Female Primary & Secondary Syphilis Incidence Rates, 1980–2017




Note: Congenital syphilis rates are per 100,000 live births; female P&S rates are per 100,000 female population.




Early Syphilis* Incidence among Females of Childbearing Age (15-44) & Congenital Syphilis Incidence by County, California, 2017

Female Early Syphilis Rates




PublicHealth

Congenital Syphilis Rates



^{*} Includes primary, secondary, and early latent syphilis.


Congenital Syphilis Cases versus Female Early Syphilis* Incidence Rates, California, 2008–2017





^{*} Includes primary, secondary, and early latent syphilis.


Congenital Syphilis Cases versus Female Early Syphilis* Cases by Pregnancy Status California, 2008–2017





^{*} Includes primary, secondary, and early latent syphilis.


Congenital Syphilis Cases versus Females of Childbearing Age (15-44) Early Syphilis* Cases by Pregnancy Status California, 2008-2017





^{*} Includes primary, secondary, and early latent syphilis.


Congenital Syphilis Cases versus Female Syphilis* Cases by Pregnancy Status California, 2008-2017





^{*} Includes primary, secondary, early latent, and late latent/unknown duration syphilis.


Female Early Syphilis* and Congenital Syphilis Cases California, 2009–2017



^{*} Includes primary, secondary, and early latent syphilis.




Female Syphilis* and Congenital Syphilis Cases California, 2009–2017



^{*} Includes primary, secondary, early latent, and late latent syphilis.



Congenital Syphilis Incidence Rates by Race/Ethnicity of Mother California, 2008–2017






Note: A/PI = Asian/Pacific Islander

Data excluded for Native American/Alaskan Native due to small numbers of cases causing unstable rates.


Congenital Syphilis Incidence Rates by Race/Ethnicity of Mother California, 2017





Note: Chart excludes Native American/Alaskan Native (0 cases)


Congenital Syphilis Number of Cases by Race/Ethnicity of Mother California, 2017





Note: Chart excludes Native American/Alaskan Native (0 cases)

Congenital Syphilis Incidence Rates by Race/Ethnicity of Mother California, 2017



Note: Chart excludes Native American/Alaskan Native (0 cases) and race/ethnicity unknown (23 cases)

