

BARBARA FERRER, Ph.D., M.P.H., M.Ed.

Director

MUNTU DAVIS, M.D., M.P.H.

**Health Officer** 

CYNTHIA A. HARDING, M.P.H.

Chief Deputy Director

JEFFREY D. GUNZENHAUSER, M.D., M.P.H.

Director, Disease Control Bureau

MARIO J. PÉREZ, M.P.H.

Director, Division of HIV and STD Programs

600 South Commonwealth Avenue, 10th Floor Los Angeles, CA 90005 TEL (213) 351-8001 • FAX (213) 387-0912

www.publichealth.lacounty.gov

September 24, 2018

Dear Colleague,

OUT OF LOS ANGERED

BOARD OF SUPERVISORS

Hilda L. Solis First District

Mark Ridley-Thomas

Sheila Kuehl Third District

Janice Hahn Fourth District

Kathryn Barger Fifth District

With full support of the Los Angeles County Board of Supervisors, the Department of Public Health's Division of HIV and STD Programs (DHSP) has launched a public health education campaign to increase provider awareness of congenital syphilis and rising rates of syphilis in women.

- Increases in **female syphilis** cases led to a dramatic **increase of congenital syphilis** cases in LA County with a total 34 cases in 2016, and 44 cases to date in 2017.
- Congenital syphilis is a *preventable* condition that results from untreated syphilis during pregnancy, with potentially severe consequences for infected infants.

This **Syphilis in Women Action Kit** includes clinical resources for your practice and educational materials for your patients. The kit is structured around these *four* core syphilis prevention practices for women:

- 1. Take a thorough sexual history from all patients as part of routine medical care.
- 2. Screen all women of child-bearing age (15-44 years) for syphilis at least once.
- 3. **Screen all pregnant women** for syphilis during the first trimester or at their initial prenatal visit. **Re-screen pregnant women** for syphilis early in the third trimester (28-32 weeks) **and** at delivery.
- 4. *Stage syphilis correctly in order to treat correctly.* Assess for signs and symptoms of syphilis in last 12 months to determine if early or late syphilis.

For help interpreting test results and guidance on appropriate staging and timely treatment, call the Clinical Guidance & Nursing Unit consultation line at 213-368-7441.

We look forward to partnering with you to implement these new syphilis prevention strategies.

Mario J. Pérez, MPH

Director, Division of HIV and STD Programs

Leo Moore, MD, MSPHM

Acting Medical Director, Division of HIV and STD Programs